


# (12) UK Patent Application (19) GB (11) 2 036 780 A

(21) Application No 7940712  
 (22) Date of filing 26 Nov 1979  
 (30) Priority data  
 (31) 2851787  
 (32) 30 Nov 1978  
 (33) Fed. Rep. of Germany (DE)  
 (43) Application published  
 2 Jul 1980  
 (51) INT CL<sup>3</sup>  
 C09B 43/16 D06P 1/39  
 (52) Domestic classification  
 C4P 116 2G6B 2H10  
 2H11 2H15 2H20 2H21  
 2H3 2H5 2H6 2H9 9A7B  
 (56) Documents cited  
 GB 760347  
 (58) Field of search  
 C4P  
 (71) Applicants  
 Bayer Aktiengesellschaft,  
 509 Leverkusen-  
 Bayerwerk, Germany,  
 Fed. Rep. of Germany  
 (72) Inventors  
 Klaus Kunde, Karl Heinz  
 Schündehütte, Peter Wild  
 (74) Agents  
 Carpmaels & Ransford

(54) Disazo dyestuffs

(57) Dyestuffs of the formula




in which

R<sub>1</sub> is H, halogen, alkoxy or acylamino,  
 R<sub>2</sub> and R<sub>3</sub> are independently H, halogen, alkyl, alkoxy, or acylamino,  
 R<sub>4</sub> is halogen, OR<sub>5</sub>, SR<sub>5</sub> or NR<sub>6</sub>R<sub>7</sub>, wherein  
 R<sub>5</sub> is H, C<sub>1</sub>—C<sub>6</sub>-alkyl, aryl, heteroaryl, aralkyl, or cycloalkyl, and  
 R<sub>6</sub> and R<sub>7</sub> are independently H, amino, optionally substituted C<sub>1</sub>—C<sub>6</sub> alkyl,  
 aryl, heteroaryl, aralkyl or cycloalkyl, or together may complete a 5- or 6-  
 membered hetero ring optionally containing further hetero atoms, and wherein  
 rings A and B may be optionally substituted and their use for the dyeing of  
 cellulose containing materials in bright yellow shades.

GB 2 036 780 A

# SPECIFICATION Disazo dyestuffs

The invention relates to new yellow disazo dyestuffs of the general formula I



5 in which


$R_1$  denotes hydrogen, halogen, alkoxy or acylamino,  
 $R_2$  and  $R_3$  independently of one another denote hydrogen, halogen, alkyl, alkoxy or acylamino and  
 $R_4$  denotes halogen,  $OR_5$ ,  $SR_5$  or  $NR_6R_7$ ,

wherein

10  $R_5$  denotes hydrogen, low-molecular alkyl, aryl, hetaryl, aralkyl or cycloalkyl and  
 $R_6$  and  $R_7$  independently of one another denote hydrogen, amino, low-molecular alkyl, aryl, hetaryl,  
 aralkyl or cycloalkyl, or together form a 5-membered or 6-membered ring, optionally with the  
 inclusion of one or more hetero-atoms, in particular oxygen and/or nitrogen, it being possible for  
 alkyl, aryl, hetaryl and aralkyl to be in turn substituted,

15 and in which

the rings A and B can optionally be substituted, and to a process for their preparation,  
 characterised in that one mol of a cyanuric trihalide, one mol of an aminoazo compound of the general  
 formula II



20 in which

$R_1$ ,  $R_2$  and  $R_3$  have the abovementioned meaning one mol of an aminoazo compound of the  
 general formula III



in which


25 the rings A and B can optionally be substituted, and optionally one mol of a compound of the  
 general formula IV



in which

30  $R_4$  is other than halogen,  
 are reacted with one another in any desired sequence in the presence of an acid-binding agent, the  
 aminoazo compounds of the general formulae II and III being prepared in a known manner.

Advantageous dyestuffs are those of the general formula V



in which

35  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  have the abovementioned meaning and o, p, q, r and s are 0 or 1, but  
 $o + p + q + r + s$  equals 2,

35


and in which

the rings A and B can optionally be substituted.

Particularly advantageous dyestuffs are those of the general formulae I and V

in which


- 5  $R_4$ , o, p, q, r and s have the abovementioned meaning and 5  
 $R_1$  denotes  $\text{NHCOCH}_3$  or  $\text{NHCONH}_2$  when  
 $R_2$  and  $R_3$  represent H, or  
 $R_2$  denotes H,  $\text{CH}_3$ ,  $\text{OCH}_3$  or  $\text{OC}_2\text{H}_5$  when  
 $R_1$  and  $R_3$  represent H,
- 10 and in which 10  
the rings A and B can optionally be substituted.  
Very particularly advantageous dyestuffs are those of the general formulae I and V
- in which
- 15  $R_1, R_2, R_3, o, p, q, r$  and  $s$  have the abovementioned meaning, 15  
 $R_4$  denotes OH,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$ ,  $\text{NH}_2$ ,  $\text{NHC}_2\text{H}_4\text{OH}$ ,  $\text{N}(\text{C}_2\text{H}_4\text{OH})_2$ ,  $\text{NC}_2\text{H}_4\text{OC}_2\text{H}_4$ ,  $\text{NHC}_6\text{H}_5$ ,  
 $\text{N}(\text{CH}_3)\text{C}_6\text{H}_5$ ,  $\text{NH}(3-\text{C}_6\text{H}_4\text{SO}_3\text{H})$  or  $\text{NH}(4-\text{C}_6\text{H}_4\text{SO}_3\text{H})$ ,  $\text{N}(\text{CH}_3)\text{C}_2\text{H}_4\text{OH}$ ,  $\text{N}(\text{C}_2\text{H}_5)\text{C}_2\text{H}_4\text{OH}$ ,  
 $\text{NHCH}_2\text{CH}(\text{CH}_3)\text{OH}$  or  $\text{N}(\text{CH}_2\text{CH}(\text{CH}_3)\text{OH})_2$  and  
the rings A and B can optionally be substituted.  
Interesting dyestuffs are those of the general formulae I and V
- 20 in which 20  
 $R_1, R_2, R_3, o, p, q, r$  and  $s$  have the abovementioned meaning,  
 $R_4$  denotes  $\text{NH}_2$ ,  $\text{NHC}_2\text{H}_4\text{OH}$ ,  $\text{N}(\text{C}_2\text{H}_4\text{OH})_2$  or  $\text{N}(\text{CH}_3)\text{C}_2\text{H}_4\text{OH}$
- and
- 25 the rings A and B can optionally be substituted. 25  
Dyestuffs of the general formulae I and B which are prepared using compounds of the general  
formula VI



VI

in which


- 30  $R_6$  and  $R_9$  independently of one another denote H, Cl, OH,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$ ,  $\text{OCOCH}_3$ ,  $\text{OCOC}_6\text{H}_5$ , 30  
 $\text{OSO}_2\text{CH}_3$ ,  $\text{OSO}_2\text{C}_6\text{H}_5$ ,  $\text{CH}_3$ ,  $\text{CH}_2\text{SO}_3\text{H}$ ,  $\text{NH}_2$ ,  $\text{NHCOCH}_3$ ,  $\text{NHCOCH}_2\text{OH}$ ,  $\text{NHCOC}_6\text{H}_5$ ,  $\text{NHCONH}_2$ ,  
 $\text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{C}_6\text{H}_5$ ,  $\text{COOH}$ , or  $\text{SO}_3\text{H}$ ,
- and in which
- the ring B can optionally be substituted,
- as compounds of the general formula III are preferred.
- 35 Dyestuffs of the general formulae I and V which are prepared using compounds of the general 35  
formula VII



VII

in which

- 40  $R_{10}$  denotes H,  $\text{CH}_3$ , Cl,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$  or  $\text{SO}_3\text{H}$  and 40  
 $R_{11}$  denotes H,  $\text{CH}_3$ , Cl,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$ ,  $\text{NHCOCH}_3$  or  $\text{NHCONH}_2$ ,
- and in which
- the ring B can optionally be substituted,
- as compounds of the general formula III are particularly preferred.
- Dyestuffs of the general formulae I and V which are prepared using compounds of the general
- 45 formula VIII 45



VIII

in which

$R_{12}$  denotes H, Cl,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$  or OH and

$S_1$  and  $S_2$  independently of one another denote H,  $\text{SO}_3\text{H}$ ,  $\text{COOH}$  or  $\text{CH}_2\text{SO}_3\text{H}$ ,  
and in which

the ring A can optionally be substituted,  
as compounds of the general formula III are very particularly preferred.

5 Particularly interesting dyestuffs are those of the general formulae I and V which are prepared  
using compounds of the general formula VIII

5

in which

$R_{12}$  denotes 4—H, 4— $\text{CH}_3$ , 4—OH or 4— $\text{OCH}_3$ , when

$S_1$  represents 3— $\text{SO}_3\text{H}$ , 3— $\text{COOH}$  or 3— $\text{CH}_2\text{SO}_3\text{H}$  and

10  $S_2$  represents H, or

10

$R_{12}$  denotes 3—H, when

$S_1$  represents 4— $\text{SO}_3\text{H}$ , 4— $\text{COOH}$  or 4— $\text{CH}_2\text{SO}_3\text{H}$  and

$S_2$  represents H,


and in which

15 the ring A can optionally be substituted,

15

as compounds of the general formula III.

Dyestuffs of the general formula IX



in which

20 o, p, q, r and s are 0 or 1, but o + p + q + r + s equals 2,

20

and in which

$R_1$  denotes  $\text{NHCOCH}_3$  or  $\text{NHCONH}_2$ , when

$R_2$  and  $R_3$  represent H, or

$R_2$  denotes H,  $\text{OCH}_3$  or  $\text{OC}_2\text{H}_5$ , when

25  $R_1$  and  $R_3$  represent H, and

25

$R_4$  denotes  $\text{NH}_2$ ,  $\text{NHC}_2\text{H}_4\text{OH}$ ,  $\text{N}(\text{C}_2\text{H}_4\text{OH})_2$  or  $\text{N}(\text{CH}_3)\text{C}_2\text{H}_4\text{OH}$ ,

$R_{10}$  denotes H,  $\text{CH}_3$ , Cl,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$  or  $\text{SO}_3\text{H}$ ,

$R_{11}$  denotes H,  $\text{CH}_3$ , Cl,  $\text{OCH}_3$ ,  $\text{OC}_2\text{H}_5$ ,  $\text{NHCOCH}_3$  or  $\text{NHCONH}_2$  and

$R_{12}$  denotes 4—H, 4— $\text{CH}_3$ , 4—OH or 4— $\text{OCH}_3$ , when

30 S represents 3— $\text{SO}_3\text{H}$ , 3— $\text{COOH}$  or 3— $\text{CH}_2\text{SO}_3\text{H}$ , or

30

$R_{12}$  denotes 3—H, when

S represents 4— $\text{SO}_3\text{H}$ , 4— $\text{COOH}$  or 4— $\text{CH}_2\text{SO}_3\text{H}$ ,

are of very particular interest.

Examples of aminoazo compounds of the general formula II are those which are obtained by

35 diazotising 2-aminonaphthalenedisulphonic acids, such as, for example, 2-aminonaphthalene-1,5-  
disulphonic acid, 2-aminonaphthalene-3,6-disulphonic acid, 2-aminonaphthalene-3,7-disulphonic acid,  
2-aminonaphthalene-4,8-disulphonic acid, 2-aminonaphthalene-5,7-disulphonic acid or 2-  
aminonaphthalene-6,8-disulphonic acid, and coupling the diazotisation products to aminobenzenes  
which are optionally substituted in the 2-position or in the 3-position, such as, for example, aniline, 2-  
40 methylaniline, 2-methoxyaniline, 2-ethoxyaniline, 2-chloroaniline, 3-acetylaniline or 3-  
aminophenylurea; the  $\omega$ -methanesulphonic acids of the abovementioned aminobenzenes can equally  
well be used as coupling components, the protective group being split off again, by treatment with acids  
or alkalis, after the coupling.

40

Examples of aminoazo compounds of the general formula III are those which are obtained by

45 diazotising optionally substituted aminobenzenes, such as, for example, aniline, 2-chloroaniline, 3-  
chloroaniline, 4-chloroaniline, 2-methylaniline, 3-methylaniline, 4-methylaniline, 2-methoxyaniline, 3-  
methoxyaniline, 4-methoxyaniline, 2-ethoxyaniline, 3-ethoxyaniline, 4-ethoxyaniline, 2-  
aminobenzenesulphonic acid, 3-aminobenzenesulphonic acid, 4-aminobenzenesulphonic acid,  
anthranilic acid, 3-aminobenzoic acid, 4-aminobenzoic acid, 5-amino-2-hydroxybenzoic acid, 5-amino-  
50 2-methylbenzenesulphonic acid, 3-aminophenylmethanesulphonic acid or 4-  
aminophenylmethanesulphonic acid, and coupling the diazotisation products to optionally substituted  
aminobenzenes, such as, for example, aniline, 2-methylaniline, 3-methylaniline, 2-methoxyaniline, 3-  
methoxyaniline, 2-ethoxyaniline, 3-ethoxyaniline, 2-methoxy-5-methylaniline, 2-ethoxy-5-  
methylaniline, 5-methoxy-2-methylaniline, 2-aminophenol, 3-aminophenol, 2,5-dimethylaniline, 3,5-  
55 dimethylaniline, 2,5-dimethoxyaniline, 3-acetylaminoaniline, 3-methylsulphonylaminoaniline, 3-  
aminophenylurea, 1,3-diaminobenzene, 3-amino-4-methylbenzenesulphonic acid, 3-amino-4-  
hydroxybenzenesulphonic acid, 3-amino-4-methoxybenzenesulphonic acid or 3-

45

50

55

aminophenylmethanesulphonic acid; the  $\omega$ -methanesulphonic acids of the abovementioned aminobenzenes can equally well be used as coupling components, the protective group being split off again, by treatment with acids or alkalis, after the coupling.

5 Examples of aminoazo compounds of the general formula III are also those which are obtained by diazotising optionally further substituted 4-nitroaminobenzenes, such as, for example, 4-nitroaniline, 4-nitro-2-methoxyaniline, 4-nitroaniline-2-sulphonic acid or 4-nitro-5-methoxyaniline-2-sulphonic acid, and coupling the diazotisation products to optionally substituted hydroxybenzenes, such as, for example, phenol, salicylic acid, 3-hydroxybenzoic acid, 2-hydroxy-6-methylbenzoic acid, 2-hydroxybenzenesulphonic acid or 3-hydroxybenzenesulphonic acid, and then reducing the nitroazo compound to give the aminoazo compound. 10

Examples of aminoazo compounds of the general formula III are also those which are obtained on treating other aminoazo compounds of the general formula III with agents with which sulphonic acid radicals can be introduced, such as, for example, sulphuric acid, oleum or sulphur trioxide, an example being 4-amino-3,4'-azobenzenedisulphonic acid.

15 Examples of compounds of the general formula IV are water, methanol, ethanol, glycol, 2-methoxyethanol, phenol, thiophenol, ammonia, hydrazine, methylamine, ethylamine, dimethylamine, diethylamine, ethanolamine, diethanolamine, N-methylethanolamine, N-ethylethanolamine, glycine, N-methylglycine, taurine, N-methyltaurine, aminomethanesulphonic acid, N-methylaminomethanesulphonic acid, aniline, N-methylaniline, 3-aminobenzenesulphonic acid, 4-aminobenzenesulphonic acid, 2-aminonaphthalene-4,8-disulphonic acid, 2-aminopyridine, 2-aminothiazole, benzylamine, pyrrolidine, piperidine, morpholine, 1-amino-2-propanol and bis-(2-hydroxypropyl)-amine. 20

The reaction between the cyanuric halide, for example cyanuric fluoride, cyanuric chloride or cyanuric bromide, the aminoazo compounds of the general formulae II and III and the compounds of the general formula IV is carried out in three stages in any desired sequence, the first stage being carried out at about 0—10°, the second at about 35—50° and the third at about 80—110°C, and the acid thereby formed being neutralised with alkaline agents, such as, for example, sodium acetate, sodium bicarbonate, sodium carbonate, sodium hydroxide solution, lithium carbonate, lithium hydroxide, potassium carbonate or potassium hydroxide. 25

Those aminoazo compounds or compounds of the general formula IV which carry groups conferring solubility in water are advantageously reacted first. 30

The dyestuffs are precipitated from the solution by adding salt and are isolated and dried or isolated by spray-drying. In general, they are obtained in the form of salts, in particular the alkali metal salts, and preferably the sodium salts. The formulae given are those of the free acids.

35 The new dyestuffs dye cellulose-containing materials in clear yellow colour shades. 35

In the examples which follow, "parts" denote parts by weight, "percentages" denote percentages by weight and the temperature is given in degrees Centigrade.

#### EXAMPLE 1

29.5 parts of the aminoazo compound, of the general formula II, obtained from 2-aminonaphthalene-4,8-disulphonic acid and aniline are dissolved in 1,000 parts of water, and a solution of 13.5 parts of cyanuric chloride in 100 parts of acetone is added at 0°, the hydrochloric acid liberated being neutralised by adding 46.5 parts of an aqueous 20% strength solution of sodium carbonate. When the first reaction stage has ended, 22.2 parts of the aminoazo compound, of the general formula III, obtained from 3-aminobenzenesulphonic acid and 2-methoxyaniline are dissolved in 1,000 parts of water and the solution is added to the first mixture. The temperature is increased to 40° and the hydrochloric acid liberated is neutralised by adding 46.5 parts of 20% strength sodium carbonate solution. When the second reaction stage has ended, 15.2 parts of diethanolamine are added. The temperature is increased to 90°, and the mixture is then stirred at this temperature for a further 3 hours. The dyestuff is precipitated by adding 450 parts of potassium chloride and is isolated. After drying, an orange-coloured powder, an aqueous solution of which dyes cellulose-containing materials in greenish-tinted yellow shades, is obtained. 50

#### EXAMPLES 2 to 4

Very similar dyestuffs are obtained if, instead of the diethanolamine used in Example 1, 18 parts of an aqueous 25% strength solution of ammonia or 8.9 parts of ethanolamine or N-methylethanolamine are used. 55

#### EXAMPLE 5

18.6 parts of the aminoazo compound, of the general formula III, which is obtained by diazotising 4-nitroaniline, coupling the diazotisation product to salicylic acid and then reducing the coupling product are dissolved in 1,000 parts of water and the solution is reacted with a suspension of 13.5 parts of cyanuric chloride in 100 parts of water at 0°, the hydrochloric acid liberated being neutralised by adding 46.5 parts of an aqueous 20% strength solution of sodium carbonate. When the first reaction stage has ended, 29.5 parts of the aminoazo compound, of the general formula II, which is obtained 60

Example	Compound of the general formula III, obtained from	Compound of the general formula IV
66	4-aminophenylmethanesulphonic acid and 3-acetylaminoaniline	ammonia
67	3-aminophenylmethanesulphonic acid and 3-acetylaminoaniline	diethanolamine
68	"	ethanolamine
69	"	ammonia
70	4-aminobenzenesulphonic acid and 3-aminophenylurea	diethanolamine
71	"	ethanolamine
72	"	ammonia
73	3-aminobenzenesulphonic acid and 3-aminophenylurea	diethanolamine
74	"	ethanolamine
75	"	ammonia
76	4-aminobenzoic acid and 3-aminophenylurea	diethanolamine
77	"	ethanolamine
78	"	ammonia
79	3-aminobenzoic acid and 3-aminophenylurea	diethanolamine
80	"	ethanolamine
81	"	ammonia
82	5-amino-2-methylbenzenesulphonic acid and 3-aminophenylurea	diethanolamine
83	"	ethanolamine
84	"	ammonia
85	4-aminophenylmethanesulphonic acid and 3-aminophenylurea	diethanolamine
86	"	ethanolamine
87	"	ammonia
88	3-aminophenylmethanesulphonic acid and 3-aminophenylurea	diethanolamine
89	"	ethanolamine
90	"	ammonia
91	4-nitroaniline and salicylic acid	diethanolamine
92	"	ethanolamine
93	"	ammonia

Example	Compound of the general formula III, obtained from	Compound of the general formula IV
38	3-aminobenzoic acid and 2-methoxyaniline	ethanolamine
39	"	ammonia
40	5-amino-2-methylbenzenesulphonic acid and 2-methoxyaniline	diethanolamine
41	"	ethanolamine
42	"	ammonia
43	4-aminophenylmethanesulphonic acid and 2-methoxyaniline	diethanolamine
44	"	ethanolamine
45	"	ammonia
46	3-aminophenylmethanesulphonic acid and 2-methoxyaniline	diethanolamine
47	"	ethanolamine
48	"	ammonia
49	4-aminobenzenesulphonic acid and 3-acetylaminoaniline	diethanolamine
50	"	ethanolamine
51	"	ammonia
52	3-aminobenzenesulphonic acid and 3-acetylaminoaniline	diethanolamine
53	"	ethanolamine
54	"	ammonia
55	4-aminobenzoic acid and 3-acetylaminoaniline	diethanolamine
56	"	ethanolamine
57	"	ammonia
58	3-aminobenzoic acid and 3-acetylaminoaniline	diethanolamine
59	"	ethanolamine
60	"	ammonia
61	5-amino-2-methylbenzenesulphonic acid and 3-acetylaminoaniline	diethanolamine
62	"	ethanolamine
63	"	ammonia
64	4-aminophenylmethanesulphonic acid and 3-acetylaminoaniline	diethanolamine
65	"	ethanolamine

Example	Compound of the general formula III, obtained from	Compound of the general formula IV
7	4-aminobenzenesulphonic acid and aniline	diethanolamine
8	"	ethanolamine
9	"	ammonia
10	3-aminobenzenesulphonic acid and aniline	diethanolamine
11	"	ethanolamine
12	"	ammonia
13	4-aminobenzoic acid and aniline	diethanolamine
14	"	ethanolamine
15	"	ammonia
16	3-aminobenzoic acid and aniline	diethanolamine
17	"	ethanolamine
18	"	ammonia
19	5-amino-2-methylbenzenesulphonic acid and aniline	diethanolamine
20	"	ethanolamine
21	"	ammonia
22	4-aminophenylmethanesulphonic acid and aniline	diethanolamine
23	"	ethanolamine
24	"	ammonia
25	3-aminophenylmethanesulphonic acid and aniline	diethanolamine
26	"	ethanolamine
27	"	ammonia
28	4-aminobenzenesulphonic acid and 2-methoxyaniline	diethanolamine
29	"	ethanolamine
30	"	ammonia
31	3-aminobenzenesulphonic acid and 2-methoxyaniline	diethanolamine
32	"	ethanolamine
33	"	ammonia
34	4-aminobenzoic acid and 2-methoxyaniline	diethanolamine
35	"	ethanolamine
36	"	ammonia
37	3-aminobenzoic acid and 2-methoxyaniline	diethanolamine



from 2-aminonaphthalene-4,8-disulphonic acid and aniline are dissolved in 1,000 parts of water and the solution is added to the first mixture. The temperature is increased to 40° and the subsequent procedure is as in Example 1. An orange-coloured powder, an aqueous solution of which dyes cellulose-containing materials in greenish-tinged yellow shades, is obtained.

5 EXAMPLE 6

Very similar dyestuffs are obtained if, instead of the diethanolamine used in Example 5, 18 parts of 5 an aqueous 25% strength solution of ammonia, 8.9 parts of ethanolamine or 10.9 parts of N-methylethanolamine are used.

EXAMPLES 7 to 96

10 If the procedure of the processes indicated in Examples 1—6 is followed and the product obtained 10 from 2-aminonaphthalene-4,8-disulphonic acid and aniline is used as the compound of the general formula II and the compounds indicated in the table below are used as the compounds of the general formulae III and IV, yellow dyestuffs for cellulose-containing materials are likewise obtained:


Example	Compound of the general formula III, obtained from	Compound of the general formula IV
94	4-aminoazobenzene and 2 mols of SO <sub>3</sub> (4-aminoazobenzene-2,4'-disulphonic acid)	diethanolamine
95	"	ethanolamine
96	"	ammonia

Similarly good dyestuffs are obtained using N-methylethanolamine as compound IV.

Similarly good dyestuffs are obtained if, instead of the compound of the general formula II used in Examples 1—96, those obtained from 2-aminonaphthalene-4,8-disulphonic acid and 2-methoxyaniline or 3-acetaminoaniline and 3-aminophenylurea are used.

## 5 CLAIMS

### 1. A dyestuff of the general formula



in which


- 10  $R_1$  denotes a hydrogen or halogen atom or an alkoxy or acylamino group,  
 $R_2$  and  $R_3$  independently of each other denote a hydrogen or halogen atom or an alkyl, alkoxy, or  
acylamino group and  
 $R_4$  denotes a halogen atom or an OR<sub>5</sub>, SR<sub>5</sub> or NR<sub>6</sub>R<sub>7</sub> group

in which

- 15  $R_5$  denotes a hydrogen atom, C<sub>1</sub> to C<sub>6</sub> alkyl, aryl, hetaryl, aralkyl or cycloalkyl and  
 $R_6$  and  $R_7$  independently of each other denote a hydrogen atom or an amino, C<sub>1</sub> to C<sub>6</sub> alkyl, aryl,  
hetaryl, aralkyl or cycloalkyl group, or together form a 5-membered or 6-membered ring,  
optionally with the inclusion of one or more hetero-atoms, and the alkyl, aryl, hetaryl and aralkyl  
groups are optionally substituted,

and in which

- 20 the rings A and B are optionally substituted.  
2. A dyestuff according to claim 1 in which  $R_6$  and  $R_7$  together form a 5-membered or 6-  
membered ring with the inclusion of one or more oxygen and/or nitrogen atoms.  
3. A dyestuff according to claim of the general formula



- 25 in which  
 $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  have the same meanings as in claim 1 and  
 $o$ ,  $p$ ,  $q$ ,  $r$  and  $s$  are 0 or 1, but  $o + p + q + r + s$  equals 2,

and in which

- the rings A and B are optionally substituted.  
30 4. A dyestuff according to any of the foregoing claims in which  $R_4$  has the same meaning as in  
claim,

- $o$ ,  $p$ ,  $q$ ,  $r$  and  $s$  have the same meaning as in claim 3 and  
 $R_1$  denotes NHCOCH<sub>3</sub> or NHCONH<sub>2</sub> when  
 $R_2$  and  $R_3$  denote hydrogen atoms, or  
35  $R_2$  denotes a hydrogen atom or a methyl, methoxy, or ethoxy group when

$R_1$  and  $R_3$  denote hydrogen atoms,  
and in which

the rings A and B are optionally substituted.

5. A dyestuff according to any of the foregoing claims in which

5  $R_1$ ,  $R_2$  and  $R_3$  have the same meanings as in claim 1 or 4,

$o$ ,  $p$ ,  $q$ ,  $r$  and  $s$  have the same meaning as in claim 3 and

$R_4$  denotes OH, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>4</sub>OCH<sub>3</sub>, NH<sub>2</sub>, NHC<sub>2</sub>H<sub>4</sub>OH, N(C<sub>2</sub>H<sub>4</sub>OH)<sub>2</sub>, N(CH<sub>3</sub>)C<sub>2</sub>H<sub>4</sub>OH, N(C<sub>2</sub>H<sub>5</sub>)C<sub>2</sub>H<sub>4</sub>OH, NHCH<sub>2</sub>CH(CH<sub>3</sub>)OH, N(CH<sub>2</sub>CH(CH<sub>3</sub>)OH)<sub>2</sub>, N(C<sub>2</sub>H<sub>4</sub>OC<sub>2</sub>H<sub>4</sub>), NHC<sub>6</sub>H<sub>5</sub>, N(CH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>, NH(3—C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>H) or NH(4—C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>H)

10 and in which

the rings A and B are optionally substituted.

6. A dyestuff according to claim 5,

in which

$R_1$ ,  $R_2$  and  $R_3$  have the same meaning as in claim 1 or 4,

15  $o$ ,  $p$ ,  $q$ ,  $r$  and  $s$  have the same meaning as in claim 3, and


$R_4$  denotes NH<sub>2</sub>, NHC<sub>2</sub>H<sub>4</sub>OH, N(C<sub>2</sub>H<sub>4</sub>OH)<sub>2</sub> or N(CH<sub>3</sub>)C<sub>2</sub>H<sub>4</sub>OH,

and in which

the rings A and B are optionally substituted.

7. A dyestuff according to any of the foregoing claims in which the moiety consisting of rings A

20 and B and the intervening azo bridge is a radical of the general formula



(Va)

in which

$R_8$  and  $R_9$  independently of each other denote a hydrogen or chlorine atom or an OH, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub>,


OCOCCH<sub>3</sub>, OCOC<sub>6</sub>H<sub>5</sub>, OSO<sub>2</sub>CH<sub>3</sub>, OSO<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, CH<sub>3</sub>, CH<sub>2</sub>SO<sub>3</sub>H, NH<sub>2</sub>, NHCOCH<sub>3</sub>, NHCOCH<sub>2</sub>OH,

25 NHCOC<sub>6</sub>H<sub>5</sub>, NHCONH<sub>2</sub>, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, COOH or SO<sub>3</sub>H group,

and in which

the ring B is optionally substituted.

8. A dyestuff according to claim 7 in which the radical (Va) is of the general formula



(Vb)

30 in which

$R_{10}$  denotes H, CH<sub>3</sub>, Cl, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub> or SO<sub>3</sub>H and


$R_{11}$  denotes H, CH<sub>3</sub>, Cl, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub>, NHCOCH<sub>3</sub> or NHCONH<sub>2</sub>,

and in which

the ring B is optionally substituted.

35 9. A dyestuff according to any of the foregoing claims in which the moiety consisting of rings A

and B and the intervening azo bridge is a radical of the general formula



(Vc)

in which

$R_{12}$  denotes a hydrogen or chlorine atom or a methyl, methoxy, ethoxy or hydroxyl group

40  $S_1$  and  $S_2$  independently of each other denote a hydrogen atom or a SO<sub>3</sub>H, COOH or CH<sub>2</sub>SO<sub>3</sub>H group,

and in which

the ring A is optionally substituted.

10. A dyestuff according to claim 9 in which

45  $R_{12}$  denotes 4—H, 4—CH<sub>3</sub>, 4—OH or 4—OCH<sub>3</sub>, when

$S_1$  denotes 3—SO<sub>3</sub>H, 3—COOH or 3—CH<sub>2</sub>SO<sub>3</sub>H and

45


$S_2$  denotes a hydrogen atom,  
 $R_{12}$  denotes 3—H, when  
 $S_1$  denotes 4—SO<sub>3</sub>H, 4—COOH or 4—CH<sub>2</sub>SO<sub>3</sub>H and  
 $S_2$  denotes a hydrogen atom.

5 and in which

the ring A can optionally be substituted.

11. A dyestuff according to claim 1 of the general formula

5



in which

10  $o, p, q, r$  and  $s$  are 0 or 1, but  $o + p + q + r + s$  equals 2,

and in which

$R_1, R_2$  and  $R_3$  have the same meanings as in claim 4,

$R_4$  has the same meaning as in claim 6,

$R_{10}$  and  $R_{11}$  have the same meanings as in claim 8, and


15  $R_{12}$  and  $S$  have the same meanings as in claim 10.

12. A dyestuff according to claim 1 as hereinbefore specifically identified.

13. A process for the production of a dyestuff as claimed in claim 1 in which, per mol a cyanuric trihalide, one mol of an aminoazo compound of the general formula

10


15



20 in which

$R_1, R_2$  and  $R_3$  have the same meanings as in claim 1, one mol of an aminoazo compound of the general formula

20



in which

25 the rings A and B are optionally substituted, if  $R_4$  is other than a halogen atom, one mol of a compound of the general formula

25



in which

30  $R_4$  has the same meaning as in claim 1, other than a halogen atom, are reacted with one another in any desired sequence in the presence of an acid-binding agent.

30

14. A process for the production of a dyestuff as claimed in claim 1 when carried out substantially as described in any one of the Examples.

15. A dyestuff as claimed in claim 1 when produced by the process of claim 14.

35 16. A process for dyeing a cellulose-containing material comprising treating the material with a dyestuff as claimed in any of claims 1 to 12 and 15.

35

17. A cellulose-containing material when dyed by the process of claim 6.